

Online Supplementary Document

Wu et al. Socioeconomic status and prevalence of type 2 diabetes in mainland China, Hong Kong and Taiwan: a systematic review

J Glob Health 2017;7:011103

Appendix S1. PRISMA Checklist

Section/topic	#	Checklist item	Reported
TITLE			
Title	1	Identify the report as a systematic review, meta-analysis, or both.	Yes
ABSTRACT			
Structured summary	2	Provide a structured summary including, as applicable: background; objectives; data sources; study eligibility criteria, participants, and interventions; study appraisal and synthesis methods; results; limitations; conclusions and implications of key findings; systematic review registration number.	Yes
INTRODUCTION			
Rationale	3	Describe the rationale for the review in the context of what is already known.	Yes
Objectives	4	Provide an explicit statement of questions being addressed with reference to participants, interventions, comparisons, outcomes, and study design (PICOS).	Yes
METHODS			
Protocol and registration	5	Indicate if a review protocol exists, if and where it can be accessed (e.g., Web address), and, if available, provide registration information including registration number.	Yes

Eligibility criteria	6	Specify study characteristics (e.g., PICOS, length of follow-up) and report characteristics (e.g., years considered, language, publication status) used as criteria for eligibility, giving rationale.	Yes
Information sources	7	Describe all information sources (e.g., databases with dates of coverage, contact with study authors to identify additional studies) in the search and date last searched.	Yes
Search	8	Present full electronic search strategy for at least one database, including any limits used, such that it could be repeated.	Yes
Study selection	9	State the process for selecting studies (i.e., screening, eligibility, included in systematic review, and, if applicable, included in the meta-analysis).	Yes
Data collection process	10	Describe method of data extraction from reports (e.g., piloted forms, independently, in duplicate) and any processes for obtaining and confirming data from investigators.	Yes
Data items	11	List and define all variables for which data were sought (e.g., PICOS, funding sources) and any assumptions and simplifications made.	Yes
Risk of bias in individual studies	12	Describe methods used for assessing risk of bias of individual studies (including specification of whether this was done at the study or outcome level), and how this information is to be used in any data synthesis.	Yes
Summary measures	13	State the principal summary measures (e.g., risk ratio, difference in means).	Yes
Synthesis of results	14	Describe the methods of handling data and combining results of studies, if done, including measures of consistency (e.g., I^2) for each meta-analysis.	Yes
Risk of bias across studies	15	Specify any assessment of risk of bias that may affect the cumulative evidence (e.g., publication bias, selective reporting within studies).	No
Additional analyses	16	Describe methods of additional analyses (e.g., sensitivity or subgroup analyses, meta-regression), if done, indicating which were pre-specified.	No
RESULTS			
Study selection	17	Give numbers of studies screened, assessed for eligibility, and included in the review, with reasons for exclusions at each stage, ideally with a flow diagram.	Yes
Study characteristics	18	For each study, present characteristics for which data were extracted (e.g., study size, PICOS, follow-up period) and provide the citations.	Yes
Risk of bias within studies	19	Present data on risk of bias of each study and, if available, any outcome level assessment (see item 12).	Yes

Results of individual studies	20	For all outcomes considered (benefits or harms), present, for each study: (a) simple summary data for each intervention group (b) effect estimates and confidence intervals, ideally with a forest plot.	Yes
Synthesis of results	21	Present results of each meta-analysis done, including confidence intervals and measures of consistency.	Yes
Risk of bias across studies	22	Present results of any assessment of risk of bias across studies (see Item 15).	No
Additional analysis	23	Give results of additional analyses, if done (e.g., sensitivity or subgroup analyses, meta-regression [see Item 16]).	No
DISCUSSION			
Summary of evidence	24	Summarize the main findings including the strength of evidence for each main outcome; consider their relevance to key groups (e.g., healthcare providers, users, and policy makers).	Yes
Limitations	25	Discuss limitations at study and outcome level (e.g., risk of bias), and at review-level (e.g., incomplete retrieval of identified research, reporting bias).	Yes
Conclusions	26	Provide a general interpretation of the results in the context of other evidence, and implications for future research.	Yes
FUNDING			
Funding	27	Describe sources of funding for the systematic review and other support (e.g., supply of data); role of funders for the systematic review.	Yes

From: Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group (2009). Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 6(7): e1000097. doi:10.1371/journal.pmed1000097

For more information, visit: www.prisma-statement.org.

Appendix S2. Systematic review search terms

Medline (Ovid, 1946 to May Week 1 2016)

- 1 socioeconomic status.mp. or Social Class/
- 2 Socioeconomic Factors/ or Social Class/ or social position.mp.
- 3 Income/ or income.mp.
- 4 Education/ or education.mp.
- 5 occupation.mp. or Occupations/
- 6 1 or 2 or 3 or 4 or 5
- 7 China.mp. or China/
- 8 hongkong.mp or Hong Kong/
- 9 Taiwan.mp or Taiwan/
- 10 7 or 8 or 9
- 11 diabetes.mp. or Diabetes Mellitus, Type 2/ or Diabetes Mellitus/
- 12 6 and 10 and 11

Embase (1980 to 2016 Week 19)

- 1 socioeconomic status.mp. or social status/
- 2 social class.mp. or social class/
- 3 social status/ or social class/ or social position.mp. or socioeconomics/
- 4 income.mp. or income/
- 5 education.mp. or education/
- 6 occupation/ or occupation.mp.
- 7 1 or 2 or 3 or 4 or 5 or 6
- 8 China.mp. or China/
- 9 hongkong.mp or Hong Kong/
- 10 taiwan.mp. or Taiwan/
- 11 8 or 9 or 10
- 12 diabetes.mp. or diabetes mellitus/
- 13 7 and 11 and 12

Global Health (1973 to 2016 Week 17)

- 1 socioeconomic status.mp. or socioeconomic status/
- 2 social class.mp. or social classes/
- 3 social position.mp. or socioeconomic status.sh. or social classes.sh.
- 4 socioeconomic factors.mp. or socioeconomic status.sh. or socioeconomics.sh.
- 5 income/ or income.mp.
- 6 education/ or education.mp.
- 7 occupation.mp. or occupations/
- 8 1 or 2 or 3 or 4 or 5 or 6 or 7
- 9 china.mp. or China/
- 10 hongkong.mp or China.gl. or Hong Kong.gl.
- 11 Taiwan/ or Taiwan.mp.
- 12 9 or 10 or 11
- 13 diabetes.mp. or diabetes/ or diabetes mellitus/ or type 2 diabetes/

14 8 and 12 and 13

Appendix S3. Criteria list for the assessment of study quality based on modified Newcastle-Ottawa Scale for cohort studies

Selection: (Maximum 3 scores)

1) Representativeness of the sample:

- a) Truly representative of the average in the target population (all subjects or random sampling or from linkage database) (1 score)
- b) Somewhat representative of the average in the target population (non-random sampling)
- c) Selected group of users
- d) No description of the sampling strategy

2) Description of sample:

- a) Study year, sample size, and age clearly described (1 score)
- b) No description or not clearly described

3) Ascertainment of the exposure:

- a) The ascertainment of SES exposure clearly described (1 score)
- b) No description or not clearly described

Comparability: (Maximum 2 scores)

- 1) Confounding factors are controlled (No paper reporting both standardized prevalence and adjusted odds ratio)
 - a) Report both crude prevalence and adjusted odds ratio (2 scores)
 - b) Report standardized prevalence or adjusted odds ratio (1 score)
 - c) Report either of crude prevalence or crude odds ratio

Outcome: (Maximum 3 scores)

1) Assessment of the outcome:

- a) Laboratory test (2 scores)
- b) Record linkage (2 scores)
- c) Self report (1 score)
- d) No description

2) Statistical test:

- a) Report confidence intervals or the probability level (p value) for prevalence or odds ratio of type 2 diabetes for population with different SES (1 score)
- b) Neither confidence intervals nor the probability level (p value) are given.

Table S4: Findings of studies identified in systematic review reporting association between prevalence of type 2 diabetes and education in mainland China, Hong Kong and Taiwan

Study	Education	Adjustment		Prevalence (95% CI, %)			P-value	
				Total	Men	Women		
Wu et al. 2016 [51]	Highest level completed	2006 China population age standardized	Primary or below: Secondary: Matriculation or above:	4.1 (3.7,4.5) 4.8 (4.2,5.4) 4.6 (4.1,5.1)	4.3 (3.7,4.9) 5.5 (4.6,6.4) 5.8 (5.1,6.5)	3.9 (3.4,4.4) 4.2 (3.4,5.0) 3.1 (2.6,3.6)	p<0.05 ^a in all groups; p>0.05 ^b in all groups	
Yu et al. 2015 [74]	Highest level completed	6 th China national population census, Jilin province population standardized (standardized for various factors)	Junior school: Junior high school: High school: Undergraduate:	12.6 (11.6,13.7) 7.4 (6.7,8.3) 7.9 (7.0,8.9) 4.7 (3.9,5.6)			p<0.001 ^a	
Xu et al. 2015 [52]	School years completed	6 th China national population census age standardized	0 years: 1-9 years: ≥ 10 years:	5.4 9.2 8.8	6.6 9.1 9.8	4.5 11.9 11.6	p<0.05 ^b in total population and men	
Hu et al. 2009 [63]	Highest level completed	2000 China population census standardized (standardized for various factors)	Less than high school: High school or higher:	5.55 5.28			p>0.05 ^a	
Cai et al. 2013 [40]	Highest level completed	Crude	Illiterate: Primary school: Middle school or higher:	9.2 (8.2,10.2) 6.1 (5.2,7.0) 4.5 (3.8,5.2)			p<0.01 ^a , CI calculated from raw data	
Shi et al. 2011 [41]	Highest level completed	Crude	Primary: Junior school: High school: University:	2.36 (1.55,3.16) 3.09 (2.04,4.15) 3.16 (1.40,4.92) 3.90 (0.8,25)			p=0.616 ^a , CI calculated from raw data	
Lin et al. 2011 [42]	Highest level completed	Crude	Elementary school and below: Junior and senior high school: College/university and above:	19.61 (16.35,22.87) 11.10 (9.10,13.09) 8.19 (6.29,10.08)			p<0.001 ^a , CI calculated from raw data	
Liu et al. 2016 [38]	School years completed	Crude	0-6 years: >7 years:	21.8 (19.2,24.4) 21.1 (18.9,23.3)	10.9 (8.3,13.4) 30.1 (27.8,32.4)		No p-value, CI calculated from raw data	
Bragg et al. 2014 [35]	Highest level completed	Age and study area structure of the study population standardized	No formal schooling: Primary school: Middle school: High school: College or university:	2.8 2.7 3.3 3.5 4.1	3.4 3.6 4.0 3.8	4.1 2.9 2.8 2.9 2.7	3.8 3.2 2.8 2.4 2.0	No p-value
Xia et al. 2013 [39]	Highest level completed	Crude	Lower than junior high school: Junior high school and above:	5.70 (5.13,6.27) 4.84 (4.29,5.40)			No p-value, CI calculated from raw data	
Woo et al. 2003 [44]	Highest level completed	Crude	Primary and lower: Secondary:	9.32 (6.09,12.56) 4.60 (2.84,6.37)			No p-value, CI calculated from raw data	

			Post-secondary:	3.76 (0.51,7.00)	
Chou and Chi. 2005 [43]	Highest level completed	Crude	None or some elementary school:	14.08 (12.22,15.95)	p>0.05 ^a , CI calculated from raw data
			Elementary graduates or above:	11.93 (9.05,14.82)	

^a p-value for Chi-square test.

^b p-value for trend Chi-square test.

Table S5: Findings of studies identified in systematic review reporting odds ratio of type 2 diabetes for different education groups in mainland China, Hong Kong and Taiwan

Study	Education	Adjustment	Odds ratio (95% CI)			P-value
Xue et al. 2015 [75]	Highest level completed	Age, BMI, sex, marital status, income, alcohol consumption, smoking, cardiometabolic parameters and uric acid levels	Per level:	0.915 (0.785,1.066)		p=0.255
Wang et al. 2013 [77]	Highest level completed	Age, sex, yearly household income, ethnicity, BMI, current smoking, current drinking, hypertensive, township mean yearly income, township percent primary education or higher, township population size, township percentage of ethnic minorities	Illiteracy: Primary or above:	1.00 0.95 (0.91,0.99)		p=0.027
Kavikondala et al. 2011 [36]	Highest level completed	Age, sex, study phase, WHR, BMI, maternal literacy by sex, maternal literacy by Childhood SEP and maternal literacy by late adult SEP	Junior middle or lower: Senior middle or higher:	1.00 0.98 (0.78, 1.22)		No p-value
Ning et al. 2009 [53]	School years completed	Age, waist circumference, family history of diabetes, hypertension, personal income, socioeconomic status, occupation, current smoking, current drinking, leisure time physical activity, survey years and resident districts	Urban areas Men Women ≤9 years: 10-12 years: ≥13 years: Rural areas Men Women ≤9 years: 10-12 years: ≥13 years:	1.00 0.97 (0.73,1.28) 0.80 (0.62,1.03) 0.79 (0.56,1.10) 0.92 (0.60,1.42) 1.00 0.82 (0.53,1.26) 0.92 (0.57,1.50) 1.00 0.82 (0.53,1.26) 0.92 (0.60,1.42) 0.92 (0.60,1.42) 0.48 (0.16,1.47)		No p-value
Chen et al. 2001 [49]	Highest level completed	Age, sex, Fukiens, Hakaas, Aborigines, family history, BMI, employment, single, smoker, drinker, regular exercise, cholesterol, triglyceride and hypertension	Illiterate: Literate:	1.00 0.72 (0.47,1.11)		p>0.05
Zhou et al. 2015 [55]	Highest level completed	Region, urban/SEC cross classification, sex/age cross classification, health literacy, time since health check, minor psychiatric morbidity, weight status, number of unhealthy lifestyles, economic status, couple status and living alone	None: Primary: Secondary: University:	1.00 1.00 (0.95,1.05) 0.93 (0.87,0.99) 0.92 (0.83,1.01)		p>0.05 p<0.05 p>0.05
Bu et al. 2015 [76]	Highest level completed	Sex, family history of diabetes, TG per increase of 0.56mM (50mg/dl), BMI, systolic blood pressure per income of 10 mmHg, age per 10-yr increment	Elementary school: High school: Tertiary or above:	1.52 (1.08,2.14) 1.42 (1.13,1.80) 1.00		p<0.001 p<0.001

Zhang et al. 2013 [28]	School years completed	Age, sex, cigarette smoking, alcohol drinking, physical inactivity, BMI, hypertension, and family history of diabetes	≤ 6 years: 7-9 years: > 9 years:	Undiagnosed 0.73(0.51,1.03) 0.82 (0.59,1.16) 1.00	Diagnosed 1.11 (0.80,1.53) 0.95 (0.70,1.28) 1.00	No p-value
Cai et al. 2011 [29]	Highest level completed	Age, sex, current smokers, current drinkers, hypertension, overweight, central obesity, family history of diabetes	Illiteracy: Primary: Middle or higher:	1.00 0.81 (0.65,0.96) 0.73 (0.52,0.93)		p<0.05 p<0.05
Tai et al. 1992 [50]	Highest level completed	Age, sex, family income, exercise, physical activity, family history of diabetes and BMI	Illiterate: Elementary and junior high school: Senior high school:	1.00 1.01 0.93		p>0.05 p>0.05
Chou et al. 1994 [45]	Highest level completed	Crude	Primary school and illiterate: Junior high and above:	1.00 0.39 (0.24,0.62)		No p-value

BMI - body mass index, WHR - waist-hip ratio, SEP - socioeconomic position, SEC - socioeconomic circumstances, TG - triglyceride

Table S6: Findings of studies identified in systematic review reporting both prevalence and odds ratio of type 2 diabetes for different education groups in mainland China, Hong Kong and Taiwan

Study	Education	Outcome	P-value
Fu et al. 2011 [30]	School years completed	Crude prevalence (95% CI, %)	No p-value, CI calculated from raw data
		≤ 9 years: > 9 years:	
		Odds ratio (95% CI) adjusted for age, sex, BMI, smoking alcohol drinking, and regular leisure physical activity	p=0.787
		≤ 9 years: > 9 years:	
Wei et al. 2010 [78]	Highest level completed	Crude prevalence (95% CI, %)	No p-value, CI calculated from raw data
		Illiterate: Semiliterate: Primary: Secondary: College:	
		12.00 (4.60,19.40) 3.23 (0,9.55) 6.50 (2.12,10.88) 7.73 (5.49,9.98) 5.24 (2.66,7.83)	

		Odds ratio (95% CI) adjusted for age, gender, nationality, smoking history, alcohol consumption, marital status and family history of diabetes	Illiterate: 1.00 Semiliterate: 0.25 (0.03,2.08) Primary: 0.57 (0.20,1.58) Secondary: 0.93 (0.39,2.23) College: 0.62 (0.23,1.70)		p=0.1989 p=0.2779 p=0.8683 p=0.3521
Zhou et al. 2009 [37]	Highest level completed	Crude prevalence (95% CI, %)		Men Women	No p-value, CI calculated from raw data
			Elementary-school/illiterate: Secondary school: College:	23.14 (15.59,30.69) 31.19 (27.24,35.14) 22.55 (18.92,26.18) 16.15 (14.13,18.18) 31.82 (23.84,39.79) 10.42 (6.54,14.29)	
		Odds ratio (95% CI) adjusted for age and area	Elementary-school/illiterate: Secondary school: College:	1.00 0.67 (0.43,1.03) 0.52 (0.28,0.96)	Women 1.00 1.30 (0.82, 2.06) 1.52 (0.91, 2.54)
Xu et al. 2006 [31]	School years completed	Crude prevalence (95% CI, %)	0-9 years: 10-12 years: ≥13 years	1.52 (1.35,1.69) 1.86 (1.50,2.22) 3.80 (3.21,4.39)	No p-value, CI calculated from raw data
		Odds ratio (95% CI) adjusted for age, gender, area of residence, BMI, occupation, smoking status, occupational physical activity, hospital category and health-care payment option	0-9 years: 10-12 years: ≥13 years:	1.00 0.76 (0.59,0.99) 0.79 (0.62,1.02)	No p-value

BMI - body mass index

Table S7: Findings of studies identified in systematic review reporting association between prevalence of type 2 diabetes and income in mainland China, Hong Kong and Taiwan

Study	Income	Adjustment	Prevalence (95% CI, %)				P-value	
Yu et al. 2015 [74]	Family monthly income (¥)	6 th China national population census, Jilin Province population standardized (standardized for various factors)	<500:	11.2 (10.0,12.4)			p<0.01 ^a	
			500-:	9.5 (8.4,10.7)				
			1000-:	8.1 (7.3,9.0)				
			2000-:	6.2 (5.4,7.3)				
			3000-:	5.8 (4.5, 7.4)				
Xu et al. 2015 [52]	Family yearly income (US\$)	6 th China national population census age standardized	<800:	Total:	Men	Women	p<0.05 ^b in all groups	
			800-1599:	4.1	5.1	2.8		
			≥1600:	5.4	5.1	5.4		
				11.3	14.2	8.8		
Hu et al. 2009 [63]	Personal yearly income (¥)	2000 China population census standardized (standardized for various factors)	<1,500:	5.9			p<0.01 ^a	
			1500-5999:	4.89				
			≥6000:	6.75				
Lin et al. 2011 [42]	Family yearly income (US\$)	Crude	< 15,000:	16.53 (14.36,18.71)			p<0.001 ^a , CI calculated from raw data	
			15,000-37,500:	8.63 (6.85,10.40)				
			> 37,500:	6.37 (3.34,9.40)				
Bragg et al. 2014 [35]	Family yearly income (¥)	Age and study area structure of the study population standardized		Self-reported		Screen-detected		No p-value
				Men	Women	Men	Women	
			< 2500:	2.4	2.9	3.7	3.9	
			2500-4999:	2.8	3.1	3.1	3.0	
			5000-9999:	2.8	3.5	2.3	2.7	
			10000-19999:	2.9	3.3	2.6	3.0	
			20000-34999:	3.1	3.4	2.7	2.8	
			≥35000:	3.8	4.1	3.4	2.7	

^a p-value for Chi-square test.

^b p-value for trend Chi-square test.

Table S8: Findings of studies identified in systematic review reporting odds ratio of type 2 diabetes for different income groups in mainland China, Hong Kong and Taiwan

Study	Income	Adjustment	Odds ratio (95% CI)				P-value
Xue et al. 2015 [75]	Family monthly income (¥)	Age, BMI, sex, marital status, education, alcohol consumption, smoking, cardiometabolic parameters and uric acid levels	Per level (<999,1000-2999, ≥3000):		0.896 (0.790,1.015)		p=0.085
Wang et al. 2013 [77]	Family yearly income (¥)	Age, sex, education level, ethnicity, BMI, current smoking, current drinking, hypertensive, township mean yearly income, township percent primary education or higher, township population size, township percentage of ethnic minorities	Per 1000¥:		0.89 (0.83,0.97)		p=0.01
Ning et al. 2009 [53]	Personal monthly income (¥)	Age, waist circumference, family history of diabetes, hypertension, school years completed, socioeconomic status, occupation, current smoking, current drinking, leisure time physical activity, survey and resident districts	Urban areas	Men	Women		No p-value
			≤600:	1.00	1.00		
			601-999:	1.29 (0.93,1.79)	0.81 (0.64,1.01)		
			1000-1999:	1.35 (0.93,1.95)	0.92 (0.66,1.27)		
			≥2000:	0.99 (0.61,1.59)	0.61 (0.36,1.42)		
			Rural areas	Men	Women		
			≤600:	1.00	1.00		
			601-999:	1.37 (0.88,2.12)	0.88 (0.60,1.30)		
			1000-1999:	1.29 (0.81,2.04)	1.15 (0.74,1.80)		
			≥2000:	2.00 (1.15,3.48)	1.17 (0.56,2.45)		
Zhang et al. 2013 [28]	Personal monthly income (¥)	Age, sex, cigarette smoking, alcohol drinking, physical inactivity, BMI, hypertension, and family history of diabetes	< 1000:	Undiagnosed	Diagnosed		No p-value
			1000-1999:	4.50 (3.07,6.61)	4.56 (3.20,6.48)		
			≥2000:	1.39 (0.95,2.04)	1.31 (0.92,1.86)		
				1.00	1.00		
Yan et al. 2012 [54]	Family yearly income (¥)	Crude		Men	Women		No p-value
				Aged 18-40	Aged ≥40	Aged 18-40	Aged ≥40
			low:	1.00	1.00	1.00	1.00
			Medium:	0.59 (0.24,1.45)	1.06 (0.77,1.46)	4.06 (0.47,34.84)	0.67 (0.49,0.93)
			High:	0.86 (0.39,1.91)	1.51 (1.12,2.04)	2.16 (0.22,20.84)	0.88 (0.65,1.17)
Cai et al. 2011 [29]	Family yearly income (US\$)	Age, sex, current smokers, current drinkers, hypertension,	< 450:	1.00			
			450-650:	1.06 (0.81,1.46)			
				1.22 (0.94,1.53)		p>0.05 p>0.05	

		overweight, central obesity, family history of diabetes	> 650:		
Pan et al. 1997 [64]	Personal yearly income (¥)	Age, sex, BMI, hypertension, family history, labour, education	<2500: 2500-5000: >5000:	1.00 NA 1.48	
Tai et al. 1992 [50]	Family monthly income (Taiwan\$)	Age, sex, educational level, exercise, physical activity, family history of diabetes and BMI	10000: 10001-20000: >20000:	1.00 1.13 0.89	P=0.0001 p<0.05 p>0.05
Yu and Wong. 2004 [48]	Family monthly income (HK\$)	Age and sex	Low (<10,000): High (≥10,000):	2.01 (1.04,3.88) 1.00	p<0.05

BMI - body mass index

Table S9: Findings of studies identified in systematic review reporting both prevalence and odds ratio of type 2 diabetes for different income groups in mainland China, Hong Kong and Taiwan

Study	Income		Outcome		P-value
Fu et al. 2011 [30]	Family income	Crude prevalence (95% CI, %)	Low: Medium: High:	3.0 (1.52,4.55) 1.8 (1.37,2.24) 2.6 (0.80,4.31)	No p-value, CI calculated from raw data
		Odds ratio (95% CI) adjusted for age, sex, BMI, smoking alcohol drinking, and regular leisure physical activity	Low: Medium: High:	1.79 (0.98,3.27) 1.00 1.55 (0.72,3.34)	p=0.058 p=0.261
Wei et al. 2010 [78]	Family yearly income (¥)	Crude prevalence (95% CI, %)	<10,000: 10,000-20,000: 20,000-30,000: ≥30,000:	12.30 (7.58,17.02) 5.61 (3.47,7.74) 7.08 (4.28,9.89) 4.00 (0.14,7.86)	No p-value, CI calculated from raw data
		Odds ratio (95% CI) adjusted for age, gender, nationality, education, smoking history, alcohol consumption, marital status and family history of diabetes	<10,000: 10,000-20,000: 20,000-30,000: ≥30,000:	1.00 0.44 (0.23,0.81) 0.60 (0.31,1.17) 0.32 (0.10,0.98)	p=0.0085 p=0.1337 p=0.0465
Zhou et al. 2009 [37]	Family yearly income (¥)	Crude prevalence (95% CI, %)		Men Women	No p-value, CI calculated from raw data
			<10,000: 10,000-30,000: >30,000:	15.51 (10.31,20.71) 23.17 (18.51,27.84) 31.97 (26.10,37.83)	19.69 (16.24,23.13) 19.03 (16.42,21.65) 18.91 (15.80,22.03)
		Odds ratio (95% CI) adjusted for age and area		Men Women	No p-value
			<10,000: 10,000-30,000: >30,000:	1.00 1.50 (0.92,2.45) 2.11 (1.25,3.57)	1.00 0.90 (0.66,1.22) 0.80 (0.57,1.14)

Wu et al. 2013 [46]	Family income quintile	Crude prevalence (95% CI, %)	Lowest: 2.9 (2.2,3.7) Second: 5.2 (4.2,6.3) Middle: 7.1 (5.7,8.7) Fourth: 7.3 (6.0,8.8) Highest: 9.3 (7.5,11.4)	No p-value
		Odds ratio (95% CI) adjusted for age, sex and urban/rural residence	Lowest: 1.00 Second: 1.9 (1.4,2.7) Middle: 2.3 (1.7,3.3) Fourth: 2.4 (1.6,3.6) Highest: 2.5 (1.7,3.6)	p<0.05 p<0.05 p<0.05 P<0.05
Xu et al. 2006 [31]	Family per capita monthly income	Crude prevalence (95% CI, %)	Lower: 0.41 (0.29,0.54) Middle: 1.39 (1.15,1.63) Higher: 3.67 (3.31,4.02)	No p-value, CI calculated from raw data
		Odds ratio (95% CI) adjusted for age, gender, area of residence, BMI, education, occupation, smoking status, occupational physical activity, hospital category and health-care payment option	Lower: 1.00 Middle: 1.95 (1.27,2.99) Higher: 2.88 (1.86,4.46)	No p-value

BMI - body mass index

Table S10: Findings of studies identified in systematic review reporting association between prevalence of type 2 diabetes and occupation in mainland China, Hong Kong and Taiwan

Study	Adjustment		Prevalence (95% CI, %)	P-value
Yu et al. 2015 [74]	6 th China national population census, Jilin Province population standardized (standardized for various factors)	Manual labour:	7.1 (6.5,7.7)	p<0.001 ^a
		Mental labour:	5.9 (5.1,6.9)	
		Other:	13 (11.8,14.2)	
Hu et al. 2009 [63]	2000 China population census standardized (standardized for various factors)	Professional:	5.95	p>0.05 ^a
		Labourer:	5.4	
		Other:	5.68	
Xia et al. 2013 [39]	Crude	Mental:	4.91 (4.26,5.57)	No p-value, CI calculated from raw data
		Non-mental:	5.51 (5.00,6.01)	

^a p-value for Chi-square test.

Table S11: Findings of studies identified in systematic review reporting odds ratio of type 2 diabetes for different occupation groups in mainland China, Hong Kong and Taiwan

Study	Adjustment	Odds ratio (95% CI)		P-value	
Kavikondala et al. 2011 [36]	Age, sex, study phase, WHR, BMI, maternal literacy by sex, maternal literacy by Childhood SEP and maternal literacy by early adult SEP	Low (agricultural worker, factory work or sales and service):	1.00	No p-value	
		High (administrator/manager, professional/technical, military/police):	0.99 (0.75, 1.29)		
Ning et al. 2009 [53]	Age, waist circumference, family history of diabetes, hypertension, personal income, School years completed, socioeconomic status, current smoking, current drinking, leisure time physical activity, survey and resident districts	Urban areas	Men	Women	No p-value
		Professional:	1.00	1.00	
		Not work outside the home:	1.07 (0.73,1.57)	1.48 (0.97,2.27)	
		Worker/farmer:	0.90 (0.61,1.31)	1.21 (0.77,1.91)	
		Rural areas	Men	Women	
		Professional:	1.00	1.00	
Not work outside the home:	0.73 (0.39,1.37)	0.66 (0.26,1.68)			
Worker/farmer:	0.65 (0.38,1.10)	0.72 (0.29,1.81)			
Chen et al. 1999 [79]	Age, sex	Unemployed:	1.30 (0.92,1.82)	p>0.05	
Employed:	1.00				
Zhou et al. 2015 [55]	Region, urban SEC cross classification, sex/age cross classification, health literacy, time since health check, minor psychiatric morbidity, weight, number of unhealthy lifestyles, highest education qualification, economic status, couple status and living alone	Employed:	1.00	p<0.05 p<0.05 p<0.05	
		Housewife/husband:	1.17 (1.08,1.28)		
		Retired:	1.34 (1.24,1.45)		
		Unemployed/student/other:	1.17 (1.10,1.24)		
Zhang et al. 2013 [28]	Age, sex, cigarette smoking, alcohol drinking, physical inactivity, BMI, hypertension, and family history of diabetes	Retirement:	Undiagnosed	Diagnosed	No p-value
		Unemployment:	2.01 (1.40,2.89)	3.02 (2.12,4.22)	
		Non-manual work:	0.83 (0.55,1.25)	1.25 (0.84,1.78)	
		Manual work:	1.05 (0.70,1.55)	1.23 (0.82,1.80)	
			1.00	1.00	

WHR - waist-hip ratio, BMI - body mass index, SEP - socioeconomic position, SEC - socioeconomic circumstances

Table S12: Findings of studies identified in systematic review reporting both prevalence and odds ratio of type 2 diabetes for different occupation groups in mainland China, Hong Kong and Taiwan

Study	Outcome	P-value
Fu et al. 2011 [30]	Crude prevalence (95% CI, %) Farmer: 2.3 (1.77,2.80) Not Farmer: 1.9 (1.15,2.65)	No p-value, CI calculated from raw data
	Odds ratio (95% CI) adjusted for age, sex, BMI, smoking alcohol drinking, and regular leisure physical activity Farmer: 1.10 (0.66,1.84) Not Farmer: 1.00	p=0.723
Zhou et al. 2009 [37]	Crude prevalence (95% CI, %) Worker/farmer: 21.20 (17.35,25.05) Other: 27.60 (21.26,33.94) Office working: 29.20 (21.56,36.84)	No p-value, CI calculated from raw data
	Odds ratio (95% CI) adjusted for age and area Men: 1.00 Women: 1.00 Worker/farmer: 1.00 Other: 1.21 (0.81,1.82) Office working: 1.35 (0.85,2.12)	No p-value
Chen and Chen. 2012 [47]	Crude prevalence (%) Professionals: 1.0 Senior officials and managers: 3.7 Clerks: 1.3 Salespersons, demonstrators and models: 3.3 Craft and related workers: 2.3 Plant and machine operators and assemblers: 3.5 Personal and protective services workers: 1.7 Elementary occupations: 3.6 Skilled agricultural and fishery workers: 3.9	No p-value
	Odds ratio (95% CI) adjusted for age, sex, cigarette use, and alcohol use Professionals: 1.00 Senior officials and managers: 1.79 (1.05,3.06) Clerks: 1.48 (0.83,2.65) Salespersons, demonstrators and models: 2.72 (1.45,5.11) Craft and related workers: 1.70 (0.94,3.07) Plant and machine operators and assemblers: 2.47 (1.38,4.39) Personal and protective services workers: 1.38 (0.71,2.69) Elementary occupations: 1.80 (0.89,3.63) Skilled agricultural and fishery workers: 1.14 (0.60,2.17)	p<0.05 p>0.05 p<0.01 p>0.05 p<0.01 p>0.05 p>0.05 p>0.05
Xu et al. 2006 [31]	Crude prevalence (95% CI, %) Blue Collar: 0.64 (0.51,0.76) Service: 1.28 (0.92,1.64) White Collar: 4.08 (3.70,4.47)	No p-value, CI calculated from raw data
	Odds ratio (95% CI) adjusted for age, gender, area of residence, BMI, education, smoking status, occupational physical activity, hospital category and health-care payment option Blue Collar: 1.00 Service: 1.47 (0.99,2.18) White Collar: 1.39 (1.05,1.85)	No p-value

BMI - body mass index