ONLINE SUPPLEMENTARY DOCUMENT

Title: Availability and readiness of healthcare facilities and their effects on under-five mortality in Bangladesh: Analysis of linked data

Authors: Md Nuruzzaman Khan, Nahida Islam Trisha, Md. Mamunur Rashid

Supplementary Table S1: Variables included in creating the health facility level factors

Name of health	Number	Variables considered in generating health facility level
facility level	of items	factors
factors		
		Supportive management for providers for child health services: availability of healthcare personnel who received
Basic management and administrative system of the nearest healthcare facility where child healthcare	7	training related to child health during the 24 months preceding the survey, availability of healthcare personnel who received training related to personal supervision during the 6 months preceding the survey, and availability of healthcare personnel who received training related to child health during the 24 months and personal supervision during the 6 months preceding the survey.
services are available		Training for child health service providers in the past 24 months: EPI/cold chain, IMCI, ARI, and diarrhoea diagnosis
		Child health services: outpatient curative care for sick children, growth monitoring, child vaccination services, and diagnosis of and/or treatment for child nutrition.
Degree of availability of		Equipment for child curative care services: child scale, length or height broad, thermometer, stethoscope, infant scale, growth chart, MUAC tape, timer.
child healthcare services at the nearest healthcare facility	35	Trained staff for child curative care services: IMCI (during the past 24 months), growth monitoring (during the past 24 months)
		Essential and priority medicine: ORS, amoxicillin syrup, suspension, or dispersible, cotrimoxazole syrup, suspension, or dispersible, paracetamol syrup or suspension, vitamin A capsules, mebendazole/albendazole, zinc tablets or syrup, ampicillin powder for injection, ceftriaxone powder for injection, gentamycin for injection, and benzathine benzylpenicillin for injection
		Infection control and laboratory diagnostic capacity: soap, running water, alcohol-based hand disinfectant, latex gloves,

		sharps container, waste receptacle, haemoglobin test, stool microscopy.
Readiness of the	10	IMIC guidelines, staff trained in IMIC, child scale, thermos-
mothers' homes		meter, growth chart, zinc tablets or syrup, ORS, amoxicillin
nearest		syrup, suspension, or dispersible, paracetamol syrup or
healthcare		suspension, meben-dazole/alben-dazole.
facility (where		
child healthcare		
services are		
available) to		
provide child		
healthcare		
services		
Average distance		Average distance on road communication from women's
on road		resided cluster to the nearest health facility.
communication		
from mothers'		
resided cluster to		
the nearest		
healthcare		
facility where		
child healthcare		
services are		
available		