

# Influence of lifestyle on suboptimal health: Insights from a national cross-sectional survey in China

Jie Wang<sup>1\*</sup>, Yinghao Wang<sup>1\*</sup>,  
Zheng Guo<sup>2,3\*</sup>, Zi Lin<sup>4</sup>,  
Xiangqian Jin<sup>1</sup>, Hui Niu<sup>4</sup>,  
Yibo Wu<sup>5†</sup>, Lihua Tang<sup>6†</sup>,  
Haifeng Hou<sup>7†</sup>

<sup>1</sup>School of Public Health, Shandong First Medical University & Shandong Academy of Medical Sciences, Jinan, China

<sup>2</sup>Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt University Medical Center, Nashville, Tennessee

<sup>3</sup>School of Medical and Health Sciences, Edith Cowan University, Perth, Australia

<sup>4</sup>Taian Maternity and Child Health Hospital, Taian, China

<sup>5</sup>School of Public Health, Peking University, Beijing, China

<sup>6</sup>Department of Blood Transfusion, The Affiliated Taian City Central Hospital of Qingdao University, Taian, China

<sup>7</sup>School of Public Health and The Second Affiliated Hospital of Shandong First Medical University, Taian, China

\*Joint first authorship.

†Joint senior authorship.

## Correspondence to:

Lihua Tang, MSc, Professor  
Department of Blood Transfusion, The Affiliated Taian City Central Hospital of Qingdao University  
29 Longtan Road, Taian 271000  
China  
lhtangtaian@qq.com

Yibo Wu, PhD  
School of Public Health, Peking University  
38 Xueyuan Road, Haidian District, Beijing 100191  
China  
wuyibo@bjmu.edu.cn

**Background** Suboptimal health status (SHS) is a non-clinical or pre-disease state between optimal/ideal health and disease. While its etiology remains unclear, lifestyle is considered one of the most important risk factors. We aimed to examine the effects of lifestyles on SHS through a nationwide survey in China.

**Methods** We conducted a cross-sectional survey in 148 cities across China between 20 June and 31 August 2022, on 30 505 participants from rural and urban communities gathered through stratified quota sampling. We measured SHS with the Short-Form Suboptimal Health Status Questionnaire (SHSQ-SF). We gathered information on participants' lifestyles (ie, smoking, alcohol consumption, breakfast habits, weekly food delivery frequency, intermittent fasting, sleep duration and physical activities) through face-to-face interview. We determined the relationship between lifestyle and SHS logistic regression analysis by based on odds ratios (ORs) and 95% confidence intervals (CIs).

**Results** We included 22 897 participants (female: 13 056, male: 9841), 12 108 (52.88%) of whom reported exposure to SHS. After adjusting for demographic characteristics, individuals who currently smoked (OR=1.165; 95% CI=1.058-1.283) and those who drank alcohol (OR=1.483; 95% CI=1.377-1.596) were at a higher risk of SHS than those who have never done either. In a dose-response way, takeaway food consumption was associated with a higher risk of SHS, while increased frequency of breakfast and mild-intensity exercise conversely reduced said risk. Individuals with shorter sleep duration had a higher risk of SHS when compared to those who slept for more than seven hours per day.

**Conclusions** We observed a relatively high prevalence of SHS across China, highlighting the importance of lifestyle in health promotion. Specifically, adopting healthy dietary habits, engaging in regular physical activity, and ensuring high-quality sleep are key in preventing SHS.

**Registration** Chinese Clinical Trial Registry (ChiCTR2200061046).

Suboptimal health status (SHS) is an intermediate or borderline condition between optimal health and illness, characterised by the emergence of health issues, general weakness, and/or low energy [1]. Individuals with SHS experience declines in vitality, physiological function, and their capacity for adaptation, which often leads to higher incidence of chronic or infectious diseases [1]. As an emerging public health concern, SHS has been receiving growing

attention from medical professionals [2]. However, due to inconsistencies in SHS measurements and the heterogeneity of targeted populations, reported rates of SHS vary from 20% to 80% [3,4]. There is currently no specific treatment for this condition, and patients suffering from it experience a reduced quality of life, have to frequently visit hospitals, and suffer additional non-medical expenses [5,6]. Several studies have shown that SHS is associated with the development of conditions such as diabetes mellitus, coronary heart disease, and stroke [7-10].

As a reversible stage preceding somatic or mental disorders, SHS is regarded as a lifestyle-related complaint associated with insufficient sleep, work- or study-related stress, physical inactivity, and an unhealthy diet [9,11,12]. According to previous studies, engaging in regular physical activity, maintaining a normal weight, quitting smoking, and abstaining from alcohol abuse can help minimise the risks of mortality from various chronic diseases and certain infectious diseases, with a particular focus on cardiovascular diseases and cancers [13,14]. Healthy lifestyles could enhance individuals' ability to maintain their energy balance, resulting in optimal metabolic function and a significant reduction in the risk of chronic diseases [15,16]. Conversely, unhealthy lifestyles have the opposite effect and are widely recognised as modifiable risk factors in disease prevention and management [17,18], having reached the scale of an epidemic and showing the potential to burden society by way of an increasing prevalence of chronic adult diseases in the future [19].

Chronic diseases develop through a long-term process which lasts several years following exposure to unhealthy lifestyles. Exploring related lifestyles may help prevent chronic diseases during their early phases. However, studies have shown that community residents in China adopt healthy lifestyles at a low or moderate level [20]. Additionally, little is known about the current SHS based on large-scale representative national samples in China, despite individual-level studies involving diverse populations, including college students, teachers, civil servants, businessmen, medical personnel, and general community residents [12]. To address this gap, we conducted a nationwide cross-sectional survey to explore the prevalence of SHS among Chinese residents and identify the influencing lifestyle factors.

## METHODS

### Study design and participants

We conducted a national population-based cross-sectional survey of 148 cities across China [21]. We initially determined sampling proportions based on population distribution data from the Seventh National Census Data of China, encompassing 23 provinces, five autonomous regions, and four municipalities, after which we used a multistage random sampling approach at the provincial, municipal, district/county, and community/village levels to gather our study sample.

We included permanent residents with Chinese nationality who were  $\geq 16$  years old and had sufficient ability to read and understand the questionnaires. We excluded individuals currently diagnosed with somatic diseases or psychiatric abnormalities, those with a history of psychiatric abnormalities, those who had taken medication within the past two weeks, and those currently involved in other clinical investigations. We also excluded questionnaires that had been completed in less than 240 seconds during the interviews, if they had inconsistent logic within the responses or incomplete information, and if there were duplicates.

We publicly recruited the interviewers from local universities in each city; they received training in sampling methods, research tools, and quality control, and their skills were subsequently assessed with a pre-defined training protocol. The Shaanxi Health Culture Research Center ethics review board approved the protocol, which we also registered in the Chinese Clinical Trial Registry (ChiCTR2200061046). All participants have signed the informed consent form.

### Measurements of demographic characteristics and lifestyles

We collected data on the participants' age, gender, body mass index (BMI), educational level, marital status, living area and household income per capita. Following the Chinese criteria of BMI [22], we categorised participants as thin ( $\text{BMI} < 18.5$ ), normal ( $\text{BMI} = 18.5\text{--}23.99$ ), overweight ( $\text{BMI} = 24\text{--}27.99$ ) and obese ( $\text{BMI} \geq 28$ ). Lifestyle behavioral variables were alcohol consumption, smoking status, breakfast habits, frequency of takeaway food consumption per week, intermittent fasting, sleep duration, and physical activity.

### Measurement of SHS

We measured SHS with the Short-Form Suboptimal Health Status Questionnaire (SHSQ-SF) [23], which had a Cronbach's  $\alpha$  coefficient of 0.902 and a split-half reliability of 0.863, as well as good validity and

reliability per the Kaiser-Meyer-Olkin (KMO) and Bartlett's tests. Following scoring using a Likert 5-point scale (range=0-4), we calculated a cumulative SHS score, categorising participants as either being in optimal health (total score <11) or suboptimal health (total score ≥11) [23].

## Statistical analysis

We presented continuous variables as means and standard deviations for normally distributed or medians and interquartile ranges (IQRs) for non-normally distributed data, and categorical variables as rates or percentages. We used Pearson  $\chi^2$  test to compare the differences in the prevalence of SHS between groups and applied multivariate logistic regression analysis to identify the association between SHS and lifestyles. We performed all analyses in SPSS, version 25.0 (IBM Corporation, Armonk, New York, USA) and R, version 4.2.1 (R Core Team, Vienna, Austria). Statistical significance was set at a two-tailed  $P < 0.05$ .




Figure 1. Flowchart of participant recruitment.

(55.44%) than men (49.49%) ( $P < 0.001$ ). We found no statistically significant differences in prevalence between urban (52.80%) and rural participants (53.13%) ( $P > 0.05$ ). Meanwhile unmarried ones had a higher prevalence (56.87%) than married ones (47.31%) ( $P < 0.001$ ). Participants classified as underweight had the highest prevalence of SHS (58.07%) compared to other categories ( $P < 0.001$ ).

## RESULTS

### Characteristic of study participants

We excluded 7608 respondents with incomplete data ( $n = 6426$ ) or the presence of somatic or psychological disorders ( $n = 1182$ ) (Figure 1), resulting in a final sample of 22 897 participants (female: 13 056, male: 9841), representing 23 provinces, five autonomous regions, and four municipalities across China. We recruited 5662 (24.73%) participants from rural and 17 235 (75.27%) from urban areas. Widowed or divorced participants comprised 1.85% of the sample, while smokers and drinkers were 14.55% and 31.45%, respectively (Table 1).

### The prevalence of SHS

The prevalence of SHS was 52.88% overall and was significantly higher among women

Table 1. Characteristics of study participants

Variables, n (%)	OHS (n = 10 789)	SHS (n = 12 108)	Total (n = 22 897)	$\chi^2$	P-value
<b>Age group in years</b>				215.67	<0.001
16-25	4795 (42.73)	6426 (57.27)	11 221 (49.01)		
26-35	2025 (50.73)	1967 (49.27)	3992 (17.43)		
36-45	1726 (50.06)	1722 (49.94)	3448 (15.06)		
46-55	1785 (50.8)	1729 (49.2)	3514 (15.35)		
56-65	458 (63.43)	264 (36.57)	722 (3.15)		
<b>Gender</b>				79.76	<0.001
Male	4971 (50.51)	4870 (49.49)	9841 (42.98)		
Female	5818 (44.56)	7238 (55.44)	13 056 (57.02)		
<b>Marital status</b>				208.49	<0.001
Unmarried	5531 (43.13)	7292 (56.87)	12 823 (56.00)		
Married	5085 (52.69)	4566 (47.31)	9651 (42.15)		
Divorced	134 (40.73)	195 (59.27)	329 (1.44)		
Widowed	39 (41.49)	55 (58.51)	94 (0.41)		
<b>Living area</b>				0.18	>0.05
Urban	8135 (47.2)	9100 (52.8)	17 235 (75.27)		
Rural	2654 (46.87)	3008 (53.13)	5662 (24.73)		

Table 1. continued

Variables, n (%)	OHS (n = 10 789)	SHS (n = 12 108)	Total (n = 22 897)	$\chi^2$	P-value
<b>BMI</b>				52.82	<0.001
Underweight (<18.5)	1578 (41.93)	2185 (58.07)	3763 (16.43)		
Normal (18.5-24)	6805 (48.3)	7285 (51.7)	14090 (61.54)		
Overweight (24-28)	1945 (48.38)	2075 (51.62)	4020 (17.56)		
Obese ( $\geq 28$ )	461 (45.02)	563 (54.98)	1024 (4.47)		
<b>Income per capita in CNY/mo</b>				58.78	<0.001
$\leq 3000$	3191 (43.9)	4078 (56.1)	7269 (31.75)		
3001-5999	4571 (49.12)	4734 (50.88)	9305 (40.64)		
6000-8999	1380 (50.36)	1360 (49.64)	2740 (11.97)		
9000-11 999	747 (46.08)	874 (53.92)	1621 (7.08)		
$\geq 12 000$	900 (45.87)	1062 (54.13)	1962 (8.56)		
<b>Education level</b>				160.56	<0.001
Primary school or below	659 (55.85)	521 (44.15)	1180 (5.15)		
Middle school	1313 (56.79)	999 (43.21)	2312 (10.10)		
High school	2764 (47.52)	3052 (52.48)	5816 (25.41)		
University/college	5630 (44.42)	7044 (55.58)	12 674 (55.35)		
Graduate school (e.g. master's/doctoral degree)	423 (46.23)	492 (53.77)	915 (3.99)		

BMI – body mass index, CNY – Chinese Yuan, SHS – suboptimal health status, OHS – optimal health status, mo – month

Prevalence rates were the highest for participants who stopped smoking (68.28%), followed by those who currently smoked (56.49%) and those who never smoked (51.93%) ( $P < 0.001$ ) (Table 2). Similarly, those who stopped drinking alcohol (62.42%) had the highest rates, followed by those who currently drank (60.12%) and those who never drank it (49.27%) ( $P < 0.001$ ). Participants who fasted intermittently had the higher prevalence rates (69.79%) than those who did not (50.84%) ( $P < 0.001$ ).

Table 2. Descriptive results of SHS-related lifestyles

Variables, n (%)	OHS (n = 10 789)	SHS (n = 12 108)	Total (n = 22 897)	$\chi^2$	P-value
<b>Smoking</b>				74.73	<0.001
Never smoking	9406 (48.07)	10 161 (51.93)	19 567 (85.85)		
Ceased smoking	177 (31.72)	381 (68.28)	558 (2.44)		
Currently smoking	1206 (43.51)	1566 (56.49)	2772 (12.11)		
<b>Alcohol consumption</b>				264.54	<0.001
Never drinking	7963 (50.73)	7733 (49.27)	15 696 (68.55)		
Ceased drinking	752 (37.58)	1249 (62.42)	2001 (8.74)		
Currently drinking	2074 (39.88)	3126 (60.12)	5200 (22.71)		
<b>Frequency breakfast per week</b>				533.72	<0.001
$\leq 2$ times	1612 (36.47)	2808 (63.53)	4420 (19.30)		
3-4 times	1425 (38.49)	2277 (61.51)	3702 (16.17)		
5-6 times	1433 (46.66)	1638 (53.34)	3071 (13.41)		
7 times	6319 (53.99)	5385 (46.01)	11 704 (51.12)		
<b>Number of takeaways per week</b>				330.45	<0.001
0 time	1604 (38.5)	2562 (61.5)	4166 (18.19)		
1 time	1896 (59.6)	1285 (40.4)	3181 (13.89)		
2 times	3853 (47.89)	4192 (52.11)	8045 (35.14)		
$\geq 3$ times	3436 (45.78)	4069 (54.22)	7505 (32.78)		
<b>Intermittent fasting</b>				316.82	<0.001
Yes	744 (30.21)	1719 (69.79)	2463 (10.76)		
No	10 045 (49.16)	10 389 (50.84)	20 434 (89.24)		
<b>Sleep duration per day</b>				691.30	<0.001
$\leq 5$ h	399 (31.87)	853 (68.13)	1252 (5.47)		
6h	1568 (34.12)	3028 (65.88)	4596 (20.07)		
7h	4363 (47.9)	4746 (52.1)	9109 (39.78)		
$> 7$ h	4459 (56.16)	3481 (43.84)	7940 (34.68)		
<b>Vigorous-intensity activity</b>				75.86	<0.001
Occasionally	8299 (47.06)	9337 (52.94)	17 636 (77.02)		
Sometimes	1347 (44.46)	1683 (55.54)	3030 (13.24)		
Often	544 (44.34)	683 (55.66)	1227 (5.36)		
Everyday	599 (59.66)	405 (40.34)	1004 (4.38)		

Table 2. continued

Variables, n (%)	OHS (n = 10 789)	SHS (n = 12 108)	Total (n = 22 897)	$\chi^2$	P-value
<b>Mild-intensity activity</b>				71.43	<0.001
Occasionally	1918 (40.02)	2875 (59.98)	4793 (20.93)		
Sometimes	1782 (41.32)	2531 (58.68)	4313 (18.83)		
Often	1717 (45.74)	2037 (54.26)	3754 (16.40)		
Everyday	5372 (53.52)	4665 (46.48)	10037 (43.84)		

SHS – suboptimal health status, OHS – optimal health status, h – hours

### Association between demographic information and SHS

The multivariate regression analysis indicated that women (OR=1.514; 95% CI=1.422-1.612) and individuals living in rural areas (OR=1.094; 95% CI=1.023-1.171) had a higher risk of SHS. Conversely, being married (OR=0.858; 95% CI=0.770-0.956) and having a normal BMI (OR=0.863; 95% CI=0.799-0.932) had a protective effect on health status. Additionally, income and age were found to have potential impacts on the development of SHS (Table 3).

Table 3. Logistic regression analysis of SHS-related lifestyles

Variables	Univariate model		Multivariate model	
	OR (95% CI)	P-value	OR (95% CI)	P-value
<b>Smoking</b>				
Never smoking	ref		ref	
Ceased smoking	1.993 (1.664-2.387)	<0.001	1.959 (1.613-2.380)	<0.001
Currently smoking	1.202 (1.109-1.302)	<0.001	1.165 (1.058-1.283)	<0.01
<b>Alcohol consumption</b>				
Never drinking	ref		ref	
Ceased drinking	1.710 (1.554-1.882)	<0.001	1.641 (1.479-1.820)	<0.001
Currently drinking	1.552 (1.456-1.654)	<0.001	1.483 (1.377-1.596)	<0.001
<b>Frequency of breakfast per week</b>				
≤2 times	ref		ref	
3-4 times	0.917 (0.838-1.004)	>0.05	0.910 (0.827-1.001)	>0.05
5-6 times	0.656 (0.598-0.721)	<0.001	0.702 (0.636-0.776)	<0.001
7 times	0.489 (0.456-0.525)	<0.001	0.617 (0.570-0.667)	<0.001
<b>Number of takeaways per week</b>				
0 time	ref		ref	
1 time	1.605 (1.477-1.745)	<0.001	1.367 (1.251-1.494)	<0.001
2 times	1.747 (1.606-1.901)	<0.001	1.426 (1.299-1.565)	<0.001
≥3 times	2.357 (2.144-2.590)	<0.001	1.781 (1.604-1.979)	<0.001
<b>Intermittent fasting</b>				
No	ref		ref	
Yes	2.234 (2.041-2.445)	<0.001	1.629 (1.481-1.793)	<0.001
<b>Sleep duration per day</b>				
>7h	ref		ref	
7h	1.393 (1.312-1.480)	<0.001	1.426 (1.339-1.520)	<0.001
6h	2.474 (2.294-2.667)	<0.001	2.336 (2.159-2.528)	<0.001
≤5h	2.738 (2.412-3.109)	<0.001	2.207 (1.931-2.522)	<0.001
<b>Vigorous-intensity activity</b>				
Occasionally	ref		ref	
Sometimes	1.111 (1.028-1.200)	<0.01	1.086 (1.000-1.179)	>0.05
Often	1.116 (0.993-1.254)	>0.05	1.097 (0.970-1.241)	>0.05
Everyday	0.601 (0.528-0.684)	<0.001	0.724 (0.631-0.832)	<0.001
<b>Mild-intensity activity</b>				
Occasionally	ref		ref	
Sometimes	0.948 (0.871-1.030)	>0.05	0.969 (0.887-1.059)	>0.05
Often	0.791 (0.726-0.863)	<0.001	0.857 (0.781-0.939)	<0.01
Everyday	0.579 (0.540-0.621)	<0.001	0.683 (0.633-0.737)	<0.001

Table 3. continued

Variables	Univariate model		Multivariate model	
	OR (95% CI)	P-value	OR (95% CI)	P-value
<b>Age group in years</b>				
16-25	ref		ref	
26-35	0.725 (0.674-0.779)	<0.001	0.850 (0.769-0.941)	<0.01
36-45	0.744 (0.690-0.804)	<0.001	0.983 (0.865-1.117)	>0.05
46-55	0.723 (0.670-0.780)	<0.001	1.035 (0.906-1.181)	>0.05
56-65	0.430 (0.368-0.503)	<0.001	0.680 (0.559-0.827)	<0.001
<b>Gender</b>				
Male	ref		ref	
Female	1.270 (1.205-1.338)	<0.001	1.514 (1.422-1.612)	<0.001
<b>Marital status</b>				
Unmarried	ref		ref	
Married	0.681 (0.646-0.718)	<0.001	0.858 (0.770-0.956)	<0.01
Divorced	1.104 (0.883-1.379)	>0.05	1.032 (0.802-1.330)	>0.05
Widowed	1.070 (0.709-1.615)	>0.05	1.221 (0.780-1.910)	>0.05
<b>Living area</b>				
Urban	ref		ref	
Rural	1.013 (0.954-1.076)	>0.05	1.094 (1.023-1.171)	<0.01
<b>BMI</b>				
Underweight (<18.5)	ref		ref	
Normal (18.5-24)	0.773 (0.719-0.831)	<0.001	0.863 (0.799-0.932)	<0.001
Overweight (24-28)	0.770 (0.704-0.843)	<0.001	0.935 (0.847-1.032)	>0.05
Obese (≥28)	0.882 (0.767-1.014)	>0.05	0.992 (0.854-1.152)	>0.05
<b>Income per capita in CNY/mo</b>				
≤3000	ref		ref	
≤3000	0.810 (0.762-0.862)	<0.001	0.834 (0.780-0.892)	<0.001
3001-5999	0.771 (0.706-0.842)	<0.001	0.757 (0.687-0.833)	<0.001
6000-8999	0.916 (0.822-1.020)	>0.05	0.883 (0.786-0.992)	<0.05
9000-11999	0.923 (0.835-1.021)	>0.05	0.846 (0.758-0.945)	<0.01
<b>Education level</b>				
Primary school or below	ref		ref	
Middle school	0.962 (0.836-1.108)	>0.05	0.873 (0.751-1.015)	>0.05
High school	1.397 (1.231-1.584)	<0.001	1.068 (0.927-1.230)	>0.05
University/college	1.583 (1.403-1.785)	<0.001	1.135 (0.989-1.303)	>0.05
Graduate school (e.g. master's/doctoral degree)	1.471 (1.237-1.750)	<0.001	1.203 (0.990-1.461)	>0.05


BMI – body mass index, CNY – Chinese Yuan, SHS – suboptimal health status, OHS – optimal health status, CI – confidence interval, OR – odds ratio, ref – reference, h – hours, mo – months

## Association between Lifestyles and SHS

After adjusting for age, gender, marital status, living area, household income, BMI, and education level, ceasing to smoke (OR=1.959; 95% CI=1.613-2.380) and current smoking (OR=1.165; 95% CI=1.058-1.283) compared to never smoking, as well as ceasing to drinking alcohol (OR=1.641; 95% CI=1.479-1.820) and current drinking (OR=1.483; 95% CI=1.377-1.596) compared to never drinking alcohol emerged as risk factors for SHS, as did frequency of takeaways per week (one time: OR=1.357, two times: OR=1.426, three or more times: OR=1.781), intermittent fasting (OR=1.629; 95% CI=1.481-1.793), and shorter sleep duration (five or less hours: OR=2.207, six hours: OR=2.336, six to seven hours: OR=1.426) compared to sleep duration more than seven hours (Figure 2 and Table 3).

Inversely, higher frequency of breakfast per week (five to six times: OR=0.702, seven times: OR=0.617) compared with two or fewer times per week and more frequent mild-intensity activity (often: OR=0.857; 95% CI=0.781-0.939, every day: OR=0.683; 95% CI=0.633-0.737) compared with occasional activity emerged as protective factors.






**Figure 2.** Forest map of the multivariate logistic regression analysis. CI – confidence interval, OR – odds ratio, ref – reference.

## DISCUSSION

Approximately 52.88% of the 22 897 participants aged 16–65 years in our nationwide cross-sectional study reported complaints related to SHS. Individuals exposed to unhealthy dietary habits, exercising insufficiently, smoking, drinking exercise, and sleeping less had an increased risk of SHS (**Figure 3**).



**Figure 3.** Risk and protective factors of SHS. SHS – suboptimal health status, OHS – optimal health status.

Early identification of SHS and its associated determinants (e.g. eating and activity habits), along with targeted interventions, can play a crucial role in mitigating adverse outcomes and offer a comprehensive and holistic way to prevent chronic diseases and enhance the quality of life.

The consumption of takeaway food increased over the past decade, emerging as a significant factor contributing to the elevating incidence of overweight and/or obesity due to its unfavourable nutritional contents [24-26]. This diet habit, in turn, predisposes individuals to cardiovascular diseases (CVD) [27-29], type 2 diabetes [30,31], and cancers [25].

Moreover, a previous observational study found that unhealthy diet is associated with a higher likelihood of mental health problems in the adult population [32]. Reduced consumption of well-balanced nutrient-rich foods (e.g. a lack of fruits and vegetables) and/or higher consumption of energy-dense foods was shown to be independently associated with increased levels of stress and depression [32-37]. These mental health factors are key determinants of SHS.

Individuals with regular breakfast tend to have healthier overall lifestyle patterns, a greater focus on healthy nutritional patterns, improved relationships, and increased ability to self-manage stress [38]. Moreover, habitual breakfast eaters exhibit a higher proclivity to physical activities than those with irregular breakfast [38]. We had similar findings, demonstrating a significant positive association between the prevalence of SHS and irregular breakfast consumption and the consumption of takeaway food.

Intermittent fasting leads to changes in the body's energy metabolism processes and affects the progression of various diseases. However, inconsistent with a previous study [39], we did not find it to be associated with lower risk of SHS. Moreover, the body tends to adapt to long-term regular meal times, so short-term intermittent fasting might lead to a perception of poor health, despite the long-term health benefits of more regular fasting. Physical activity and regular exercise have been shown to reduce the risk of NCDs in a dose-dependent manner, including cardiovascular diseases, type 2 diabetes, and cancers [40]. Our findings further confirm this and suggest that frequent mild-intensity physical activities are associated with a lower prevalence of SHS [41].

Sleep is an essential biological and behavioral process that significantly contributes to the maintenance of overall health and quality of life [42], acting as a modulator of cardiovascular function, blood glucose regulation, and hormonal secretion [43]. Insufficient sleep duration can impact metabolic regulation, autonomic nervous functions, blood coagulation system, and endothelial dysfunction [44]. We have observed that lack of sleep was associated with higher risk of SHS, which is in line with previous studies [45-47].

We also found a higher prevalence of SHS among smokers compared to those who have never smoked, while ex-smokers had a higher prevalence than either group. This phenomenon may be attributed to the common issue of reverse causality often encountered in cross-sectional studies, where individuals who have experienced health problems may decide to quit smoking. This is similar to the rationale behind the observation that individuals who have ceased alcohol drinking had a higher prevalence of SHS than current drinkers. Furthermore, long-term smoking and drinking often result in the body adapting to these behaviors. Therefore, individuals' cessation of these habits can disturb the body's equilibrium and potentially contribute to a perception of declining health [48].


## Limitations

We designed this study as an observational, cross-sectional survey; this allowed us to explore the national prevalence of SHS and the association between lifestyle factors and SHS, but prevented us from establishing causality, which should be the purpose of future longitudinal studies. Second, we did not define prevalence of SHS based on sex-specific cut-off points. Third, we obtained all information through self-reported questionnaires, which may introduce potential information bias. Fourthly, although we assessed SHS using a standardised questionnaire, methodological differences may exist when compared to studies conducted in other populations or employing different questionnaires. Lastly, we assessed the frequency of consuming takeaway food per week, yet the term "takeaway" encompasses a broad range of foods, some of which may be healthy, while others are not, resulting in varying effects on consumers' health.

## CONCLUSIONS

We found that unhealthy dietary habits, insufficient exercise, smoking, alcohol drinking, and short sleep duration were associated with an increased risk of SHS, underscoring the importance of adopting healthy lifestyles to enhance health status and prevent chronic diseases.





**Acknowledgments:** We would like to express our gratitude to the participants who volunteered and provided their consent for their valuable involvement in this study.

**Data availability:** The data are available from the corresponding authors on a reasonable request.

**Funding:** This work was supported by Natural Science Foundation of Shandong Province (ZR2022MH082).

**Authorship contributions:** Haifeng Hou, Yibo Wu and Lihua Tang conceived the study and guided the development of research and the preparation of manuscripts. Jie Wang, Yinghao Wang, Zheng Guo, Zi Lin, Xiangqian Jin performed the material preparation and data collection. Jie Wang, Yinghao Wang, and Zheng Guo researched data, performed the statistical analyses, and wrote the manuscript. Hui Niu, and Haifeng Hou provided critical expert advice or critical review of the current manuscript. All authors read and approved the final manuscript.

**Disclosure of interest:** The authors completed the ICMJE Disclosure of Interest Form (available upon request from the corresponding author) and disclose no relevant interests.

## REFERENCES

- 1 Wang W, Yan Y. Suboptimal health: a new health dimension for translational medicine. *Clin Transl Med*. 2012;1:28. Medline:23369267 doi:10.1186/2001-1326-1-28
- 2 Dunstan RH, Sparkes DL, Roberts TK, Crompton MJ, Gottfries J, Dascombe BJ. Development of a complex amino acid supplement, Fatigue Reviva™, for oral ingestion: initial evaluations of product concept and impact on symptoms of sub-health in a group of males. *Nutr J*. 2013;12:115. Medline:23927677 doi:10.1186/1475-2891-12-115
- 3 Bi J, Huang Y, Xiao Y, Cheng J, Li F, Wang T, et al. Association of lifestyle factors and suboptimal health status: a cross-sectional study of Chinese students. *BMJ Open*. 2014;4:e005156. Medline:24951109 doi:10.1136/bmjopen-2014-005156
- 4 Chen J, Cheng J, Liu Y, Tang Y, Sun X, Wang T, et al. Associations between breakfast eating habits and health-promoting lifestyle, suboptimal health status in Southern China: a population based, cross sectional study. *J Transl Med*. 2014;12:348. Medline:25496597 doi:10.1186/s12967-014-0348-1
- 5 Liang YZ, Chu X, Meng SJ, Zhang J, Wu LJ, Yan YX. Relationship between stress-related psychosocial work factors and suboptimal health among Chinese medical staff: a cross-sectional study. *BMJ Open*. 2018;8:e018485. Medline:29511008 doi:10.1136/bmjopen-2017-018485
- 6 Henningsen P, Zipfel S, Sattel H, Creed F. Management of Functional Somatic Syndromes and Bodily Distress. *Psychother Psychosom*. 2018;87:12-31. Medline:29306954 doi:10.1159/000484413
- 7 Adua E, Roberts P, Wang W. Incorporation of suboptimal health status as a potential risk assessment for type II diabetes mellitus: a case-control study in a Ghanaian population. *EPMA J*. 2017;8:345-55. Medline:29209438 doi:10.1007/s13167-017-0119-1
- 8 Alzain MA, Asweto CO, Zhang J, Fang H, Zhao Z, Guo X, et al. Telomere Length and Accelerated Biological Aging in the China Suboptimal Health Cohort: A Case-Control Study. *OMICS*. 2017;21:333-9. Medline:28617662 doi:10.1089/omi.2017.0050
- 9 Kupaev V, Borisov O, Marutina E, Yan YX, Wang W. Integration of suboptimal health status and endothelial dysfunction as a new aspect for risk evaluation of cardiovascular disease. *EPMA J*. 2016;7:19. Medline:27621756 doi:10.1186/s13167-016-0068-0
- 10 Hsu SH, Chen DR, Cheng Y, Su TC. Association of Psychosocial Work Hazards With Depression and Suboptimal Health in Executive Employees. *J Occup Environ Med*. 2016;58:728-36. Medline:27206130 doi:10.1097/JOM.0000000000000760
- 11 Yan YX, Dong J, Liu YQ, Yang XH, Li M, Shia G, et al. Association of suboptimal health status and cardiovascular risk factors in urban Chinese workers. *J Urban Health*. 2012;89:329-38. Medline:22203493 doi:10.1007/s11524-011-9636-8
- 12 Yan YX, Liu YQ, Li M, Hu PF, Guo AM, Yang XH, et al. Development and evaluation of a questionnaire for measuring suboptimal health status in urban Chinese. *J Epidemiol*. 2009;19:333-41. Medline:19749497 doi:10.2188/jea.JE20080086
- 13 GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2018;392:1923-94. Medline:30496105 doi:10.1016/S0140-6736(18)32225-6
- 14 Danaei G, Ding EL, Mozaffarian D, Taylor B, Rehm J, Murray CJ, et al. The preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS Med*. 2009;6:e1000058. Medline:19399161 doi:10.1371/journal.pmed.1000058
- 15 Van Cauter E, Spiegel K, Tasali E, Leproult R. Metabolic consequences of sleep and sleep loss. *Sleep Med*. 2008;9:S23-8. Medline:18929315 doi:10.1016/S1389-9457(08)70013-3
- 16 Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review. *Obesity (Silver Spring)*. 2008;16:643-53. Medline:18239586 doi:10.1038/oby.2007.118
- 17 Schulenkorf N, Siefken K. Managing sport-for-development and healthy lifestyles: The sport-for-health model. *Sport Manage Rev*. 2019;22:96-107. doi:10.1016/j.smr.2018.09.003
- 18 Nyberg ST, Singh-Manoux A, Pentti J, Madsen IEH, Sabia S, Alfredsson L, et al. Association of Healthy Lifestyle With Years Lived Without Major Chronic Diseases. *JAMA Intern Med*. 2020;180:760-8. Medline:32250383 doi:10.1001/jamainternmed.2020.0618
- 19 Chen X, Giles J, Yao Y, Yip W, Meng Q, Berkman L, et al. The path to healthy ageing in China: a Peking University-Lancet Commission. *Lancet*. 2022;400:1967-2006. Medline:36423650 doi:10.1016/S0140-6736(22)01546-X
- 20 Zhou C, Zheng W, Tan F, Lai S, Yuan Q. Influence of health promoting lifestyle on health management intentions and behaviors among Chinese residents under the integrated healthcare system. *PLoS One*. 2022;17:e0263004. Medline:35077472 doi:10.1371/journal.pone.0263004

- 21 Wang YJ, Kaierdebieke A, Fan SY, Zhang RF, Huang MJ, Li H, et al. Study protocol: A cross-sectional study on psychology and behavior investigation of Chinese residents, PBICR. *Psychosomatic medicine*. 2022;4:19. doi:10.53388/202219
- 22 Chen C, Lu FC. The guidelines for prevention and control of overweight and obesity in Chinese adults. *Biomed Environ Sci*. 2004;17 Suppl:1-36. Medline:15807475
- 23 Sun S, Liu H, Guo Z, Guan Q, Wang Y, Wang J, et al. Development and validation of a short-form suboptimal health status questionnaire. *EPMA J*. 2023. doi:10.1007/s13167-023-00339-z
- 24 Janssen HG, Davies IG, Richardson LD, Stevenson L. Determinants of takeaway and fast food consumption: a narrative review. *Nutr Res Rev*. 2018;31:16-34. Medline:29037273 doi:10.1017/S0954422417000178
- 25 Anderson AS, Key TJ, Norat T, Scoccianti C, Cecchini M, Berrino F, et al. European Code against Cancer 4th Edition: Obesity, body fatness and cancer. *Cancer Epidemiol*. 2015;39:S34-45. Medline:26205840 doi:10.1016/j.canep.2015.01.017
- 26 Duffey KJ, Gordon-Larsen P, Steffen LM, Jacobs DR Jr, Popkin BM. Regular consumption from fast food establishments relative to other restaurants is differentially associated with metabolic outcomes in young adults. *J Nutr*. 2009;139:2113-8. Medline:19776183 doi:10.3945/jn.109.109520
- 27 Smith KJ, Blizzard L, McNaughton SA, Gall SL, Dwyer T, Venn AJ. Takeaway food consumption and cardio-metabolic risk factors in young adults. *Eur J Clin Nutr*. 2012;66:577-84. Medline:22146886 doi:10.1038/ejcn.2011.202
- 28 Salas-Salvadó J, Becerra-Tomás N, García-Gavilán JF, Bulló M, Barrubés L. Mediterranean Diet and Cardiovascular Disease Prevention: What Do We Know? *Prog Cardiovasc Dis*. 2018;61:62-7. Medline:29678447 doi:10.1016/j.pcad.2018.04.006
- 29 Pallazola VA, Davis DM, Whelton SP, Cardoso R, Latina JM, Michos ED, et al. A Clinician's Guide to Healthy Eating for Cardiovascular Disease Prevention. *Mayo Clin Proc Innov Qual Outcomes*. 2019;3:251-67. Medline:31485563 doi:10.1016/j.mayocpiqo.2019.05.001
- 30 Hruby A, Manson JE, Qi L, Malik VS, Rimm EB, Sun Q, et al. Determinants and Consequences of Obesity. *Am J Public Health*. 2016;106:1656-62. Medline:27459460 doi:10.2105/AJPH.2016.303326
- 31 Hawley JA, Sassone-Corsi P, Zierath JR. Chrono-nutrition for the prevention and treatment of obesity and type 2 diabetes: from mice to men. *Diabetologia*. 2020;63:2253-9. Medline:32761356 doi:10.1007/s00125-020-05238-w
- 32 Jacka FN. Nutritional Psychiatry: Where to Next? *EBioMedicine*. 2017;17:24-9. Medline:28242200 doi:10.1016/j.ebiom.2017.02.020
- 33 Ye ZJ, Qiu HZ, Li PF, Chen P, Liang MZ, Liu ML, et al. Validation and application of the Chinese version of the 10-item Connor-Davidson Resilience Scale (CD-RISC-10) among parents of children with cancer diagnosis. *Eur J Oncol Nurs*. 2017;27:36-44. Medline:28279394 doi:10.1016/j.ejon.2017.01.004
- 34 Sarris J, Logan AC, Akbaraly TN, Paul Amminger G, Balanzá-Martínez V, Freeman MP, et al. International Society for Nutritional Psychiatry Research consensus position statement: nutritional medicine in modern psychiatry. *World Psychiatry*. 2015;14:370-1. Medline:26407799 doi:10.1002/wps.20223
- 35 Banta JE, Segovia-Siapo G, Crocker CB, Montoya D, Alhusseini N. Mental health status and dietary intake among California adults: a population-based survey. *Int J Food Sci Nutr*. 2019;70:759-70. Medline:30773065 doi:10.1080/09637486.2019.1570085
- 36 Taylor AM, Thompson SV, Edwards CG, Musaad SMA, Khan NA, Holscher HD. Associations among diet, the gastrointestinal microbiota, and negative emotional states in adults. *Nutr Neurosci*. 2020;23:983-92. Medline:30794085 doi:10.1080/1028415X.2019.1582578
- 37 Quehl R, Haines J, Lewis SP, Buchholz AC. Food and Mood: Diet Quality is Inversely Associated with Depressive Symptoms in Female University Students. *Can J Diet Pract Res*. 2017;78:124-8. Medline:28333547 doi:10.3148/cjdp-2017-007
- 38 Reeves S, Halsey LG, McMeel Y, Huber JW. Breakfast habits, beliefs and measures of health and wellbeing in a nationally representative UK sample. *Appetite*. 2013;60:51-7. Medline:23032303 doi:10.1016/j.appet.2012.09.024
- 39 Zhao X, Yang J, Huang R, Guo M, Zhou Y, Xu L. The role and its mechanism of intermittent fasting in tumors: friend or foe? *Cancer Biol Med*. 2021;18:63-73. Medline:33628585 doi:10.20892/j.issn.2095-3941.2020.0250
- 40 Dhuli K, Naureen Z, Medori MC, Fioretti F, Caruso P, Perrone MA, et al. Physical activity for health. *J Prev Med Hyg*. 2022;63:E150-E159. Medline:36479484
- 41 Eckstrom E, Neukam S, Kalin L, Wright J. Physical Activity and Healthy Aging. *Clin Geriatr Med*. 2020;36:671-83. Medline:33010902 doi:10.1016/j.cger.2020.06.009
- 42 Mosavat M, Mirsanjari M, Arabiat D, Smyth A, Whitehead L. The Role of Sleep Curtailment on Leptin Levels in Obesity and Diabetes Mellitus. *Obes Facts*. 2021;14:214-21. Medline:33756469 doi:10.1159/000514095
- 43 Knutson KL, Van Cauter E. Associations between sleep loss and increased risk of obesity and diabetes. *Ann N Y Acad Sci*. 2008;1129:287-304. Medline:18591489 doi:10.1196/annals.1417.033
- 44 Tobaldini E, Fiorelli EM, Solbiati M, Costantino G, Nobili L, Montano N. Short sleep duration and cardiometabolic risk: from pathophysiology to clinical evidence. *Nat Rev Cardiol*. 2019;16:213-24. Medline:30410106 doi:10.1038/s41569-018-0109-6
- 45 Xu T, Zhu G, Han S. Prevalence of Suboptimal Health Status and the Relationships between Suboptimal Health Status and Lifestyle Factors among Chinese Adults Using a Multi-Level Generalized Estimating Equation Model. *Int J Environ Res Public Health*. 2020;17:763. Medline:31991741 doi:10.3390/ijerph17030763
- 46 Ma C, Xu W, Zhou L, Ma S, Wang Y. Association between lifestyle factors and suboptimal health status among Chinese college freshmen: a cross-sectional study. *BMC Public Health*. 2018;18:105. Medline:29304856 doi:10.1186/s12889-017-5002-4
- 47 Tian M, Zhang Q, Zhang Z, Meng J, Liu L. Prevalence of Suboptimal Health Status and Its Influencing Factors among Chinese Software Programmers. *Iran J Public Health*. 2021;50:1361-71. Medline:34568174
- 48 Takayama S, Takase H, Tanaka T, Sugiura T, Ohte N, Dohi Y. Smoking Cessation without Educational Instruction Could Promote the Development of Metabolic Syndrome. *J Atheroscler Thromb*. 2018;25:90-7. Medline:28592705 doi:10.5551/jat.40063